VIGYAN 2047 Volume 3 No. 1 January 2026 Pages 24–26

Science behind Fasciculations in MND

Dr AK Gupta & Sanket Gupta

Muscle twitching is a familiar sensation to most people. An eyelid may flutter during stress, a calf may twitch after a strenuous run, or a random flicker may appear in the thigh while one is resting. These brief, harmless movements usually vanish as quickly as they appear. But in Amyotrophic Lateral Sclero-sis (ALS) or Motor Neuron Disease (MND), muscle twitching—known medically as fasciculations—carries a much deeper meaning. They become visible signs of what is happening inside the body at the level of its electrical wiring: the motor neurons.

To understand why fasciculations occur in ALS, it helps to think of motor neurons as power lines con-necting the brain to the muscles. These lines supply the "current" that allows muscles to contract when-ever we decide to move—whether to walk, breathe, speak, or swallow. When the power lines are intact, everything works seamlessly. But when they fray or begin to fail, the muscles they control may behave unpredictably. Just as a flickering bulb signals an unstable power supply, fasciculations are the flickers of muscle fibers receiving unstable signals from deteriorating neurons.

In ALS, the neurons primarily responsible for fasciculations are the lower motor neurons located in the anterior horn of the spinal cord and in certain regions of the brainstem. These neurons extend axons out to the muscle fibers, forming individual motor units. When ALS begins to affect these neurons, the ear-liest event is often their gradual degeneration. As individual motor neurons begin to malfunction or die, the muscle fibers they once controlled become partially denervated. Without proper electrical input, these fibers weaken, lose bulk, and become electrically unstable.

But the body does not accept this loss passively. It attempts to compensate in a remarkable way. The surviving motor neurons, sensing the loss of their neighbors, extend new branches in an effort to recon-nect with the orphaned muscle fibers. This process, called collateral sprouting, can temporarily preserve muscle function. It is somewhat like a neighborhood where many workers have quit their jobs, leaving tasks undone. The remaining workers attempt to cover multiple duties across multiple houses—not be-cause they were designed to do so, but because they want to keep things functioning. In early ALS, this process can delay noticeable weakness.

However, this overcompensation has consequences. These enlarged motor units, formed from the sur-viving neurons taking on more and more work, become electrically unstable. They cannot maintain the precise control they once had. As a result, they begin to fire spontaneously, without the brain sending a command. These spontaneous discharges of motor units appear as fasciculations, the brief rippling movements seen under the skin.



On the surface, a fasciculation may look like a tiny quiver or a brief worm-like movement. But inside the muscle, it represents a sudden burst of electrical activity from an overburdened, hyperexcitable neuron. When viewed through electromyography (EMG), these twitches appear as erratic firing patterns—fasciculation potentials—that help neurologists identify lower motor neuron dysfunction. EMG may also reveal other abnormalities such as fibrillation potentials, positive sharp waves, and signs of chronic neurogenic remodeling. These findings together support the diagnosis of ALS by showing that the motor neurons are not only dying but also attempting (and eventually failing) to compensate.

Although structural neuron degeneration explains much of the process, the biochemical environment in ALS adds another layer of complexity. Motor neurons in ALS often become hyperexcitable, meaning they fire too easily. This hyperexcitability is partly due to ion channel dysfunction. Neurons depend on sodi-um (Na^+) and potassium (K^+) channels to maintain a stable electrical state. In ALS, these channels may malfunction. Sodium channels may allow too much current inward, and potassium channels may struggle to restore the resting membrane potential. This leaves the neuron vulnerable to firing at random. An analogy can be drawn to an overcharged phone battery: one moment it jumps from 10% to 70%, the next moment it shuts off. The electrical system becomes erratic. Similarly, ALS neurons may fire spontane-ously because their internal electrical balance has become unstable.

Another biochemical contributor is glutamate excitotoxicity. Glutamate acts like the accelerator pedal in the nervous system, stimulating neurons to fire. In ALS, glutamate levels may become excessive, over-stimulating neurons and flooding them with calcium. If the accelerator is stuck and the brakes (the neu-ron's protective mechanisms) are weak, the system becomes dangerously unstable. The motor neuron fires repeatedly and uncontrollably, and over time, this leads to damage and cell death. Before the neu-ron dies, however, it may fire erratically—producing fasciculations.

Disruptions in calcium homeostasis also play a major role. Calcium acts almost like water in a dam: it must be tightly regulated. In ALS, the dam begins to leak. Calcium spills into the neuron through faulty channels or because internal buffering systems are overwhelmed. Too much calcium destabilizes the neuron's membrane and triggers spontaneous firing. Eventually, this leads to mitochondrial dysfunction and oxidative stress—further weakening the neuron. But long before the neuron dies, this biochemical chaos increases the likelihood of fasciculations.

It is important to emphasize that fasciculations do not cause weakness. They are symptoms of the motor neurons firing abnormally, not causes of muscle deterioration. Weakness in ALS arises because motor neurons are dying, not because the muscles are twitching. The twitching is just the electrical "noise" created by a failing system. A useful analogy is a damaged electrical wire: the sparks you see do not cause the wire to fail—the wire is failing, and the sparks are simply a sign of that failure. Similarly, fas-ciculations are signs of motor neuron instability, not the source of the problem.

This distinction becomes crucial when distinguishing the fasciculations of ALS from those occurring in benign conditions. The vast majority of twitching in healthy people is harmless. Stress, anxiety, exces-sive caffeine, fatigue, magnesium deficiency, and strenuous exercise can all cause twitching. These benign fasciculations are like the normal creaks a house makes at night—common, harmless, and unre-lated to structural damage. In contrast, ALS fasciculations occur alongside other alarming signs: pro-gressive muscle weakness, atrophy, abnormal EMG patterns, and spread of symptoms across different body regions. It is the combination of symptoms—not the fasciculations alone—that raises concern. Twitching by itself, especially if widespread but not accompanied by weakness or EMG abnormalities, is far more likely to be benign.

As ALS progresses, the pattern of fasciculations changes. Early in the disease, when sprouting is active and neurons are still compensating, fasciculations may be frequent and widespread. Patients may no-tice twitches in their legs, arms, face, or torso. As the disease advances, however, reinnervation be-comes insufficient. Motor neurons become overburdened and begin to die at a faster rate. Muscle fibers become permanently denervated, leading to atrophy. Fasciculations may persist, but they gradually give way to more pronounced weakness. Yet their presence remains a window into the underlying neuro-physiology of ALS.

Physiotherapy in MND

Although physiotherapy cannot stop the motor neuron degeneration underlying ALS, it plays an essential role in **preserving mobility, easing discomfort, and maintaining independence for as long as possible.** Physiotherapy in ALS is less about strengthening weakening muscles—which cannot be reversed—and more about **optimizing the function of muscles that remain innervated,** protecting joints, and preventing secondary complications such as contractures, stiffness, and pain.

A helpful way to understand physiotherapy in ALS is to imagine a **scaffolding being erected around a fragile structure.** The scaffolding does not change the structure itself, but it supports what remains, prevents collapse, and allows it to function longer and more safely. Similarly, physiotherapy provides strategic support to muscles and joints, enabling patients to move with less effort, more comfort, and reduced fatigue.

Gentle, low-impact exercises can improve **range of motion, posture, breathing mechanics, circulation, and energy management.** Techniques such as stretching, soft-tissue mobilization, and assisted movements help ease cramps and stiffness, which often accompany fasciculations and motor unit hyperexcitability. As the disease progresses, physiotherapists play a vital role in recommending **mobility aids,** orthotic supports, and adaptive equipment, helping patients conserve energy and maintain autonomy. Importantly, physiotherapy also teaches safe strategies for movement, reducing the risk of falls—a significant concern in ALS as weakness spreads.

Physiotherapy does not aim to challenge weakened muscles aggressively, because doing so can increase fatigue and sometimes accelerate discomfort. Instead, it focuses on **preservation, comfort, and energy-smart movement,** ensuring that patients live with the highest possible quality of life throughout the journey of the disease.

The Homeopathic Support

Homeopathy does not cure ALS or halt the progression of motor neuron degeneration. However, many patients find that individualized homeopathic remedies offer **comfort, emotional balance, and symptomatic relief,** particularly in managing anxiety, sleep disturbances, restlessness, cramps, and the heightened sensitivity associated with fasciculations.

Homeopathy views each patient as a unique constellation of physical and emotional responses. Rather than focusing solely on the disease label, homeopathy examines **patterns of stress, personality traits, emotional triggers, sleep quality, fatigue, cravings, and coping styles.** This individualized approach can be particularly supportive in ALS, where emotional resilience plays a significant role in overall well-being.

Certain remedies are traditionally used to address specific symptom clusters often seen in ALS:

- **Magnesia phosphorica** and **Zincum metallicum** are frequently chosen when fasciculations, cramps, and muscle twitching are distressing—similar to applying a gentle "soothing blanket" over irritable muscle fibers.
- **Gelsemium** may assist when patients experience trembling, weakness, or anticipatory anxiety.
- **Arsenicum album** is often used for restlessness, nighttime fear, and exhaustion.
- **Phosphorus** may help individuals who feel emotionally sensitive, easily fatigued, or overwhelmed by sensory stimuli.
- **Nux vomica** can support those with irritability, sleep-disturbance, and stress-driven symptoms.

The role of homeopathy in ALS is best understood through analogy: it does not repair the damaged wiring of the motor neuron system, but it can **dim the noise, cushion the emotional impact, and ease the sensory irritability** that accompanies the illness. Many patients report better sleep, reduced anxiety, greater emotional calm, and improved ability to cope with physical symptoms. ◆

__Dr AK Gupta__ is a profound homoeopathic practioner and founder director of AKGsOVIHAMS. He can be reached at drakgupta@ovihams.com

__Dr Sanket Gupta__ is a homoeopathic practioner of AKGsOVIHAMS.